

Weak Scale Quantum Gravity Effects for $\gamma\gamma \rightarrow ZZ$ in the TeV region.

S.R.Choudhury¹

Department of Physics, Delhi University, Delhi 110007, India,

A.S.Cornell² and G.C.Joshi³

*School of Physics, University of Melbourne,
Parkville, Victoria 3010, Australia*

28th of February, 2002

Abstract

We consider the effect of massive spin-2 gravitons that occur as Kaluza-Klein excitations of the graviton in a Weak scale Quantum Gravity scenario on the process $\gamma\gamma \rightarrow ZZ$, which in the Standard Model can proceed through loop diagrams. For a wide range of parameters, we show that the massive gravitons leave behind signatures that should be verifiable in a TeV scale scattering experiment.

1 Introduction

Arkani-Hamed, Dimopoulos and Dvali [1] have proposed a higher $(4 + n)$ -dimensional spacetime theory in which the Standard Model (SM) particles live in the usual 4-dimensional spacetime where as gravity also propagates in the additional compactified spatial dimensions. A remarkable feature of this scenario is the possibility that the Planck mass can become comparable to the electro-weak scale, thus solving the hierarchy mystery that naturally exists in conventional theories where the Planck mass is many orders of magnitude higher than the electro-weak scale. Gravity in our $(3+1)$ -dimensional world then becomes feeble for distances much larger than the compactification scale R . If M_{Pl} is the Planck scale in our world and M in the $(4 + n)$ -dimensional world, the two are related by [1]

$$M_{Pl}^2 = R^n M^{2+n} \quad (1)$$

Setting $M \sim \text{TeV}$, gives $R \sim \text{eV-MeV}$ for $n = 2 - 7$. Noting that the Newtonian inverse square law of gravity has been verified up to about a millimetre, [2] we see that no contradiction with Newtonian gravity results in this approach.

An alternative scenario has been proposed by Randall and Sundrum [3] where once again there exists only one fundamental Planck Scale ($\sim \text{TeV}$) with

¹src@ducos.ernet.in

²a.cornell@tauon.ph.unimelb.edu.au

³joshi@physics.unimelb.edu.au

the hierarchy of scales been generated by an exponential factor of the compactification radius in a 5-dimensional non factorizable geometry with the fifth dimension in the form of a torus. Two 3-branes with opposite tensions are assumed to reside at S_1/Z_2 orbifold fixed points $\phi = 0, \pi$ where ϕ is the angular coordinate parametrizing the extra dimension. Of the two 3-branes, it is assumed that the SM-fields reside only at the one at $\phi = \pi$ whereas gravity is all over. Calling as before M_{Pl} the 4-dimensional Planck scale, M the Planck scale in the 5-dimensional world, r_c the compactification radius of the 5-dimensional torus, the two Planck scales are related by the equation

$$M_{Pl}^2 = \frac{M^3}{k}(1 - e^{-2kr_c\pi}), \quad (2)$$

where $k \sim M_{Pl}$ is a parameter of the theory. Choosing $kr_c \sim 12$, one sees that M can have values in the TeV range, thus providing an alternative solution to the hierarchy problem.

The graviton propagating in the compact dimensions, in addition to our 4-dimensional world, appear in our world as a massless particle together with its radial excitations. In the ADD-scenario, these set of resonances are almost a continuum, each individually coupling with the 4-dimensional Planck scale strength. However because of the large number of these resonances, their combined effect mimic a weak scale spin-2 excitation. In the RS model, the graviton excitations are discrete with each coupling with masses calculable in terms of k and r_c and each coupling with TeV scale strength with matter.

Turning to phenomenological consequences of WSQG, it is clear that there will be in the nature of effective exchange effects of spin-2 particles with TeV scale coupling. Typically such contributions to the amplitudes will have suppression factor of $(\sqrt{s}/M_s)^{n+2}$ replacing the coupling parameters e or g in the corresponding SM amplitude. Clearly the most important processes where these effects will show up in the sub-TeV range are the ones where the SM-contribution is suppressed occurring only through higher order loop diagrams. Processes involving $\gamma\gamma$ collisions like $\gamma\gamma \rightarrow ZZ$ offer the most exciting possibility of such processes. The SM contributions through loops of all charged particles are well calculated and at the same time the process seems to be experimentally accessible with γ -rays obtained from backward scattering of laser beams off high energy electrons. In the present investigation, we estimate the signatures of the WSQG theories in this process for energies in the TeV-range.

2 SM-Amplitudes

The process

$$\gamma(p_1, \lambda_1) + \gamma(p_2, \lambda_2) \rightarrow Z(p_3, \lambda_3) + Z(p_4, \lambda_4) \quad (3)$$

is described by 36 invariant amplitudes $F(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4)$ where the quantities p, λ in parentheses in equation (3) denote the momentum and helicity of

the corresponding particles and s, t, u are the Mandelstam variables. The F 's are related to the corresponding differential cross-section by the relation

$$\frac{d\sigma}{d\cos\theta} = \frac{\beta_z}{64\pi s} |F|^2 \quad (4)$$

where θ is the c.m. scattering angle and $\beta_z^2 = (1 - \frac{4m_z^2}{s})$. Parity and Bose statistics imposes the following restrictions on the amplitudes

$$F(s, t, u; \lambda_1 \lambda_2 \lambda_3 \lambda_4) = F(s, t, u; \lambda_2 \lambda_1 \lambda_4 \lambda_3) (-1)^{\lambda_3 - \lambda_4} \quad (5)$$

$$F(s, t, u; \lambda_1 \lambda_2 \lambda_3 \lambda_4) = F(s, u, t; \lambda_2 \lambda_1 \lambda_3 \lambda_4) (-1)^{\lambda_3 - \lambda_4} \quad (6)$$

$$F(s, t, u; \lambda_1 \lambda_2 \lambda_3 \lambda_4) = F(s, u, t; \lambda_1 \lambda_2 \lambda_4 \lambda_3) \quad (7)$$

and

$$F(s, t, u; \lambda_1 \lambda_2 \lambda_3 \lambda_4) = F(s, t, u; -\lambda_1, -\lambda_2, -\lambda_3, -\lambda_4). \quad (8)$$

The SM amplitudes corresponding to the W-loop and the fermion loops (figure 1a) have been evaluated [4]. In the high energy limit that we are interested in, namely $s, -t, -u \gg M_{Pl}^2$ most of the amplitudes become negligible. For the W-loop contribution, the surviving amplitudes are the following together of course with their symmetry counterparts;

$$\begin{aligned} F^W(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) &= \frac{\alpha^2}{s_w^2} A^w(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) \\ A^w(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) &= \frac{(12c_w^4 - 4c_w^2 + 1)}{4c_w^2} A^s(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) \\ &\quad + \delta^w(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) \\ A^s(s, t, u; + + + +) &= 4 - \frac{4ut}{s^2} \left[\log^2 \left| \frac{t}{u} \right| + \pi^2 \right] + 4 \frac{(t-u)}{s} \log \left| \frac{t}{u} \right| \\ A^s(s, t, u; + - + -) &= 4 - \frac{4st}{u^2} \left[\log^2 \left| \frac{s}{t} \right| - 2i\pi \log \left| \frac{s}{t} \right| \right] \\ &\quad + 4 \frac{(s-t)}{u} \left[\log \left| \frac{s}{t} \right| - i\pi \right] \\ \delta^w(s, t, u; + + + +) &= 16c_w^2 \left\{ \log^2 \left| \frac{t}{u} \right| + \pi^2 + \frac{s}{u} \log \left| \frac{u}{m_w^2} \right| \log \left| \frac{s}{t} \right| \right. \\ &\quad + \frac{s}{t} \log \left| \frac{t}{m_w^2} \right| \log \left| \frac{s}{u} \right| \\ &\quad \left. - i\pi \left[\frac{s}{u} \log \left| \frac{u}{m_w^2} \right| + \frac{s}{t} \log \left| \frac{t}{m_w^2} \right| \right] \right\} \\ \delta^w(s, t, u; + - + -) &= 16c_w^2 \left\{ \log^2 \left| \frac{t}{s} \right| + \frac{u}{t} \log \left| \frac{t}{m_w^2} \right| \log \left| \frac{u}{s} \right| \right. \\ &\quad + \frac{u}{s} \log \left| \frac{s}{m_w^2} \right| \log \left| \frac{u}{t} \right| + i\pi \left[\frac{u}{t} \log \left| \frac{s}{m_w^2} \right| \right. \\ &\quad \left. \left. - \frac{u}{s} \log \left| \frac{u}{s} \right| - \frac{s^2 + t^2}{st} \log \left| \frac{t}{s} \right| \right] \right\} \quad (9) \end{aligned}$$




Figure 1a)




Figure 1b)

Figure 1: a) the SM loop graph - one single graph for all charged particles in the loop, where the dashed lines represent the Z particles. Figure 1: b) The s -channel graviton contribution graph, where the double sinusoid represents the graviton.

For the fermion loops, similarly only a few amplitudes (and their symmetric counterparts) survive as follows:

$$\begin{aligned}
 F^f(s, t, u; \lambda_1, \lambda_2, \lambda_3, \lambda_4) &= \frac{\alpha^2 Q_f^2}{4s_w^2 c_w^2} \left\{ \left[t_3^f - 2Q_f s_w^2 \right]^2 A_{\lambda_1, \lambda_2, \lambda_3, \lambda_4}^{vf}(s, t, u, m_f) \right. \\
 &\quad \left. + (t_3^f)^2 A_{\lambda_1, \lambda_2, \lambda_3, \lambda_4}^{af} \right\} \\
 A^{vf} &= -2A^s + \delta^{vf} \\
 \text{and } A^{af} &= -2A^s + \delta^{af}
 \end{aligned} \tag{10}$$

with the following nonvanishing δ 's for both the superscripts:

$$\begin{aligned}
 \delta_{++++} &= -4 \left[\log^2 \left| \frac{u}{t} \right| + \pi^2 \right] \\
 \delta_{+-+-} &= -4 \left[\log^2 \left| \frac{s}{t} \right| - i2\pi \log \left| \frac{s}{t} \right| \right]
 \end{aligned} \tag{11}$$

3 The Massive Graviton Contribution

Spin-2 massive excitations of the graviton contribute to the amplitudes via s -channel pole, as shown in fig.1(b). In the ADD version of the WSQG, the

interaction of spin-2 excitations $h_{\mu\nu}^{(n)}$ of mass m is given by the interaction Lagrangian

$$\mathcal{L}_I = -\frac{\kappa}{2} \sum_n \int d^4x T^{\mu\nu} h_{\mu\nu}^{(n)}, \quad (12)$$

where $\kappa = \sqrt{16\pi G_N}$ and $T^{\mu\nu}$ is the energy momentum tensor. Following standard Feynman rules, the contribution of fig.1(b) to the amplitudes are given by;

$$F^g(s, t, u; ++\lambda_3\lambda_4) = 0 \quad (13)$$

and the non-zero ones are;

$$\begin{aligned} F^g(s, t, u; +-++) &= D(s) \cdot \frac{sM_z^2}{2} \sin^2 \theta \\ F^g(s, t, u; +-+-) &= D(s) \cdot \frac{s^2}{2} \cos^4 \frac{\theta}{2} \\ F^g(s, t, u; +-+0) &= D(s) \cdot \frac{s^{3/2}M_z}{2\sqrt{2}} (1 + \cos \theta) \sin \theta \\ F^g(s, t, u; +- -0) &= D(s) \cdot \frac{s^{3/2}M_z}{2\sqrt{2}} (1 - \cos \theta) \sin \theta \\ F^g(s, t, u; +++-) &= D(s) \cdot \frac{sM_z^2}{2} \sin^2 \frac{\theta}{2} \\ F^g(s, t, u; +- -+) &= D(s) \cdot \frac{s^2}{2} \sin^4 \frac{\theta}{2} \\ F^g(s, t, u; +-0+) &= D(s) \cdot \frac{s^{3/2}M_z}{2\sqrt{2}} (1 - \cos \theta) \sin \theta \\ F^g(s, t, u; +-0-) &= D(s) \cdot \frac{s^{3/2}M_z}{2\sqrt{2}} (1 + \cos \theta) \sin \theta \\ F^g(s, t, u; +-00) &= D(s) \cdot \frac{s}{8} (s + 4M_z^2) \sin^2 \theta \end{aligned} \quad (14)$$

with parity symmetry yielding the rest of the amplitudes. In the last equation the quantity $D(s)$ is

$$D(s) = - \sum_n \frac{\kappa^2}{s - m_n^2}. \quad (15)$$

In the ADD scenario, these resonances almost form a continuum. The summation in the last equation can be converted into an integral and has been explicitly calculated.[5] The summation however is UV-divergent and has to be cut off at

some value M_s . M_s would be in order of magnitude of the Planck scale in $(4+n)$ -dimensions, which then is in the TeV range. Further the effective Newton constant κ in 4-dimensions would be related to M_s and the compactification size R by:

$$\kappa^2 R^n = 8\pi(4\pi)^{n/2}\Gamma(n/2)M_s^{-(n+2)}. \quad (16)$$

The result is, assuming there are n -extra compact dimensions:

$$-iD(s) = \frac{\kappa^2 s^{n/2-1} R^n}{\Gamma(n/2)} [\pi + 2iI(M_s/\sqrt{s})], \quad (17)$$

where

$$I(M_s/\sqrt{s}) = - \sum_{k=1}^{n/2-1} \frac{1}{2k} (M_s/\sqrt{s})^{2k}. \quad (18)$$

Using equation (16) , this can be written as:

$$-iD(s) = 8\pi s^{n/2-1} M_s^{-n-2} [\pi + 2iI(M_s/\sqrt{s})] \quad (19)$$

The ADD contribution is thus dependent on a single parameter M_s ($\approx TeV$) for a given n .

In the RS-version of WSQG, the interaction of the discrete set of resonances of mass m_n can be represented by

$$\mathcal{L}_I = -\frac{1}{\Lambda_\pi} \sum_n h_{\alpha\beta}^{(n)}(x) \cdot T^{\alpha\beta} \quad (20)$$

where $\Lambda_\pi = e^{kr_c\pi} M_{Pl} \approx \text{TeV}$. [6] The mass of the resonances are given in terms of the zeroes x_n of the Bessel function $J_1(x)$ by

$$m_n = kx_n e^{-kr_c\pi} \quad (21)$$

which can be written as

$$m_n = \frac{k}{M_{Pl}} m_1 \frac{x_n}{x_1}. \quad (22)$$

Phenomenologically, the RS version can be parametrized in terms of two parameters (k/M_{Pl}) which is of order 1 and m_1 , the mass of the first RS-resonance. [6] For a given value of m_1 and (k/M_{Pl}) , the summation implied in the evaluation of $D(s)$ cannot be done analytically but numerically the summation converges quite fast.

4 Results and Discussion

For the range of parameters indicated therein, our results are shown in figures 2-9. There is indeed considerable deviation from the SM results for the cross-section both in the ADD and the RS version of WSQG. The spin-2 nature of the massive gravitons is primarily responsible for the enhanced contribution of the gravitons in the TeV-range. This is responsible for the s^2 factor in their contribution to the amplitudes and combined with a dimensional coupling this starts dominating in the TeV range. In the ADD scenario, this is essentially because the net dimensionless parameter governing the amplitude is $(\sqrt{s}/M_s)^6$ for $n=2$; this is $O(1)$ in the TeV-range for $\sqrt{s}$ if M_s is in the same range.

For the RS scenario, the corresponding factor is $\left(\frac{k}{M_{Pl}}\right)^2 \frac{s}{m_1^2}$ which once again can be of $O(1)$ even for small values of the parameter (k/M_{Pl}) . The SM contribution is proportional to α^2 , so that as $\sqrt{s}$ increases, one can expect the spin-2 contribution dominating after a while. This result of course can be taken seriously only upto a point. As $\sqrt{s}/M_s$ or alternatively $(k/M_{Pl})(\sqrt{s}/m_1)$ becomes larger than one, we can no longer neglect higher order corrections that renders the amplitude unitary. Under such circumstances then, the results numerically are only indicative of considerable deviation from SM.

In conclusion, experimental data on the process $\gamma\gamma \rightarrow ZZ$ in the TeV-range and its comparison with well calculated SM - results can provide valuable signatures for possible WSQG contributions to the amplitude. It is important to note in this connection, that effects beyond the SM arising out of supersymmetric particle exchange contributions are much smaller than the WSQG effects and cannot explain if large scale departures are observed in future experimental data.




Figure 2: The Total cross-section in femto-barnes for the SM contribution and the SM plus ADD contributions for $M_s = 2, 3, 4$ TeV.




Figure 3: The Total cross-section in femto-barnes for the SM contribution and the SM plus ADD contributions for $M_s = 3, 4$ TeV.




Figure 4: The Differential cross-section in femto-barnes for the $\sqrt{s} = 0.8\text{TeV}$ against $\cos\theta$ for the SM contribution and the SM plus ADD contributions for $M_s = 2, 3, 4\text{TeV}$.




Figure 5: The Differential cross-section in femto-barnes for the $\sqrt{s} = 0.8\text{TeV}$ against $\cos\theta$ for the SM contribution and the SM plus ADD contributions for $M_s = 3, 4\text{TeV}$.




Figure 6: The Total cross-section in femto-barnes for the SM contribution and the SM plus RS contributions for $k/M_{Pl} = 0.01, 0.02, 0.03, 0.05$.




Figure 7: The Total cross-section in femto-barnes for the SM contribution and the SM plus RS contributions for $k/M_{Pl} = 0.01, 0.02, 0.03$.




Figure 8: The Differential cross-section in femto-barnes for the $\sqrt{s} = 0.8\text{TeV}$ against $\cos\theta$ for the SM contribution and the SM plus RS contributions for $k/M_{Pl} = 0.01, 0.02, 0.03, 0.05$.




Figure 9: The Differential cross-section in femto-barnes for the $\sqrt{s} = 0.8\text{TeV}$ against $\cos\theta$ for the SM contribution and the SM plus RS contributions for $k/M_{Pl} = 0.01, 0.02, 0.03$.

References

- [1] N.Arkani-Hamed, S.Dimopoulos and G.Dvali, *Phys. Lett.* **B429**, p.263 (1998).
- [2] C.Hoyle, U.Schmidt, B.Heckel, E.Adelberger, J.Gundlach, D.Kapner and H.Swanson, **hep-ph/0011014**.
- [3] L.Randall and R.Sundrum, *Phys. Rev. Lett.*, **83**, p.3370 (1999).
- [4] G.Jikia, *Nucl. Phys.* **B405**, p.24 (1993); B.Bajc, *Phys. Rev.* **D48**, p.514 (1993); M.S. Berger, *Phys. Rev.*, **D48**, p.5121 (1993); D.Dicus and C.Kao, *Phys. Rev.* **D49**, p.1265 (1994); G.Gounaris, J.Layssac, P.Porfyriadis and F.Renard, *Eur. J. Phys* **C13**, p.79 (2000).
- [5] T. Han, J.Lykken and R.Zhang, *Phys Rev.* **D59**, p.105006, (1999).
- [6] H. Davoudiasl, J.Hewett and T.Rizzo, *Phys. Rev. Lett.*, **84**, p.2080 (2000).
- [7] I.Antoniadis, N.Arkani-Hamed, S.Dimopoulos and G.Dvali, *Phys.Lett.* **B436**, p.257 (1998).